

#### Status of the AIRS-RTA



L. Strow, S. Hannon, S. De-Souza Machado, H. Motteler

- Concentrate first on the water vapor continuum
  - Fix the lower altitude channels first, then work our way up in the atmosphere
- What is the truth?
- Procedures to validate, improve the water continuum
- Error analysis, including ECMWF fields
- Fixed gas/CO<sub>2</sub> tuning, preliminary results
  - Again, truth is the problem, especially in the upper atmosphere
- Future work






## Starting Point - H<sub>2</sub>O Continuum

- January 2003 RTA quickly modified from theory based on ~1 month of validation data from Fall 2002.
- Main modifications at that time were to the water vapor continuum, primarily based on ECMWF model fields.
- Basis of changes: minimize the dependence of the bias on total column water;
  - Suggested large changes in shortwave continuum, which were implemented beyond 2400 cm<sup>-1</sup>.
  - Longwave continuum was also increased for the same reason.
- Used MT\_CKD(V1) as the starting point for validation, released Spring 2003, changes based on DOE-ARM Aeri work by U.Wisc.
- ARM-TWP has lots of water (~40 mm), and a reasonable number of "clear" overpasses.
- Goal: Agreement between AERI, AIRS-TWP, AIRS-ECMWF






#### Jan03-RTA vs Jan04-RTA??






### Jan03-RTA vs Jan04-RTA??














- Starting Point: MT\_CKD continuum released Spring 2003
- Use AIRS-TWP observations as basis for validation of RTA
  - Over ocean, good emissivity
  - Lot's of water
  - Reasonable number of "clear" observations
  - RS-90 sondes with microwave scaling for total column water
  - Determine SST with "2616 cm-1" channel, but MT\_CKD gave large biases there even though the continuum is small in this region
  - (1) Bootstrap with ECMWF-derived "2616 cm-1" bias, and/or (2) Analyze AERI shortwave data for continuum (Thanks to D. Turner and D. Tobin for the data!)
  - Once get SST, derive continuum using AIRS-TWP data
- Apply AERI, TWP continua to ECMWF, "calibrate" ECMWF statistical global column water
- Re-analyze AERI raw data using variation in radiance with total column water -MT\_CKD only used AERI data at ~40 mm total column water
- Choose "best" continuum






### Strength of Water Continuum, ARM-TWP






# ARM-TWP Biases using Continuua from 2 Analyses of AERI Data










# Lower Troposphere Water Vapor Measurement Goal: <2% in precipitable water vapor



Based on desire to improve clear sky absorption models and to resolve significant climate changes, such as the effect of CO<sub>2</sub> doubling on surface radiation budget.


For midlatitude conditions, a 10% H<sub>2</sub>O perturbation results in a ~7 W/m<sup>2</sup> change in downwelling Flux at the surface.

2% is order (1 W/m<sup>2</sup>)















#### Obs - Calc for AERI; MT\_CDK vs UMBC\_AERI








## AERI-derived Changes to MT\_CKD

















# ARM-TWP Biases for 3 Continuum Models








#### ARM-TWP for 3 Continuum Models





# UMBC

# ARM-TWP Biases for 3 Continuum Models ARM-TWP Biases vs Water Continuum Model; SST derived using 2616 cm







#### ECMWF vs TWP Biases



#### (using continuum derived from TWP)








# ECMWF Biases for 3 Continuum Models











# ECMWF Bias in Longwave - ~2% Accuracy in Total column Water?








# ECMWF Bias vs Secant( $\theta$ )








## ECMWF Bias vs Secant( $\theta$ )







# ECMWF Bias vs Secant( $\theta$ )






#### ECMWF Bias vs Total Column Water










#### AN HON

Bias for 1K ECMWF Offset above 60 mbar











- Set SST in shortwave
- Modified fixed-gas optical depth with ARM-TWP
- Strong water channels not touched
- No fixed-gas tuning above 60 mbar









## Fixed Gas Multipliers from TWP



- Trial fixed gas optical depth multipliers were generated using the ARM-TWP AIRS validation data.
- Longwave multipliers are reasonable
- 2400 cm<sup>-1</sup> multipliers seem a little large












# MT\_CKD Continuum Multipliers



- These results based on AIRS-TWP validation data in longwave, TWP and ECMWF shortwave
- Longwave multipliers agree with UMBC-AERI results below 1000 cm<sup>-1</sup>
- Still working to merge TWP and AERI in 2400 cm<sup>-1</sup> region, we are close. Linear vs quadratic?










# Sensitivity of Bias to Uniform Threshold










## Bias vs Wind Speed and Total Column Water







# ECMWF Wind Speed in Masuda $\epsilon$





#### Masuda wpeed = 0





#### Masuda wspeed = ECMWF








#### Upper Trop Water RTA Validation Remains




AWEX may help, already have good agreement between ARM-SGP, Voemel, and ECMWF (between lines)





#### ARM-SGP vs ECMWF Global Biases in H2O-sounding Channels







#### Future Work



- Pick next water continuum:
  - MT\_CKD with shortwave mods by UMBC or
  - Continuum based on AIRS co-locations with ARM-TWP sondes
  - Both models supplemented with ECMWF in extreme shortwave
  - Test with other months (Oct. 2002 done, very similar to March 2003)
  - Do high continuum values near CO<sub>2</sub> make physical sense? CO<sub>2</sub>-H<sub>2</sub>O collisions?
- Mid-wave/Upper-trop water. Awaiting ARM-CARL data and AWEX campaign results. Try our new near-wing water lineshape?
- Fixed gas modifications
  - Do RS-90's justify these mods
  - No tuning above ~60 mbar, is ECMWF biases there?
  - Any data around to test these altitudes? Limb sounders?
- Determine source of 2200 cm<sup>-1</sup> bias, N<sub>2</sub>O?



# Backup Slides









Ratio of 2 to 3 Term Fits vs Water

